

Recombinant Human DC-SIGN/CD209 Protein

Catalog No.: RP01489 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 30835 Q9NNX6-1

Tags C-His

Synonyms

CD209;CDSIGN;CLEC4L;DC-SIGN;DC-SIGN1; CLEC4L; DC-SIGN; DC-SIGN1

Product Information

Source Purification HEK293 cells ≥ 95 % as

determined by SDS-PAGE.

PAGE

Calculated MW Observed MW

40.28 kDa 40-50 kDa

Endotoxin

< 0.1 EU/ μ g of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
<u>~</u>	www.abclonal.com.cn

Background

Dendritic cell (DC)-specific intercellular adhesion molecule 3 (ICAM-3) grabbing nonintegrin (DC-SIGN), also known as CD209, is a type II transmembrane protein on DCs with a C-type lectin extracellular domain, is capable of binding ICAM-3 on resting T cells in the secondary lymphoid organs, providing the initial contact between these cells during the establishment of cell-mediated immunity. It is not only a pattern recognition receptor but implicated in immunoregulation of DCs. It has an important role in mediating DC adhesion, migration, inflammation, activating primary T cell, triggering immune response and participating in immune escape of pathogens and tumors. DC-SIGN also mediates the capture and internalization of viral, bacterial, and fungal pathogens by dendritic cells, such as HIV-1, Ebola virus, cytomegalovirus, Dengue virus, and hepatitis C virus. DC-SIGN is unique in that it regulates adhesion processes, such as DC trafficking and T-cell synapse formation, as well as antigen capture. Moreover, even though several C-type lectins have been shown to bind HIV-1, DC-SIGN does not only capture HIV-1 but also protects it in early endosomes allowing HIV-1 transport by DC to lymphoid tissues, where it enhances trans infection of T cells.

Basic Information

Description

Recombinant Human DC-SIGN/CD209 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln59-Ala404) of human CD209/DC-SIGN (Accession $\#NP_066978.1$) fused with a $6\times$ His tag at the C-terminus.

Bio-Activity

Measured by the ability of the immobilized protein to support the adhesion of ICAM-3 expressing CHO Chinese hamster ovary cells. When 5 x 104 cells/well are added to Recombinant Human DC_x001e_SIGN/CD209 Protein coated plates (5 μ g/mL with 100 μ L/well), approximately 10-20% of added cells will adhere after 1 hour at 37°C.


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human DC-SIGN/CD209 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.