

Catalog No.: RP01652 **Recombinant**

Species	Gene ID	Swiss Prot
Human	928	P21926

Synonyms
MIC3; MRP-1; BTCC-1; DRAP-27;
TSPAN29; TSPAN-29:CD9

Source	Purification
HEK293 cells	≥ 95 % as determined by SDS-PAGE

Calculated MW	Observed MW
35.64 kDa	42-47 kDa

Endotoxin
< 0.1 EU/μg of the protein by LAL method.

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

CD9 is a member of the transmembrane 4 superfamily, which is also known as the tetraspanin family. CD9 is a cell surface glycoprotein with 4 hydrophobic domains that are described as complex with integrins and other transmembrane 4 superfamily members. It is found expressed on the surface of the exosomes. The protein takes part in cellular signal transduction events and thus play a role in the regulation of cell development and activation, growth and motility. Besides, CD9 seems to be a key role in the egg-sperm fusion during the mammalian fertilization processes. CD9 is found on the membrane of the oocytes and also appears to intervene in maintaining the normal shape of oocyte microvilli.

Recombinant Human CD9 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ser112-Ile195) of human CD9 (Accession #NP_001760.1) fused with and a hFc tag at the C-terminus.


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.
After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.
Avoid repeated freeze/thaw cycles.

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Validation Data



Recombinant Human CD9 Protein was determined by SDS-PAGE under reducing (R) and non-reducing (NR) conditions.