

**Catalog No.: RP01726** **Recombinant**

Species	Gene ID	Swiss Prot
human	4914	P04629-1

C-6His

MTC; TRK; TRK1; TRKA; Trk-A; p140-TrkA; NTRK1

<b>Source</b>	<b>Purification</b>
HEK293 cells	≥ 90 % as determined by SDS-PAGE

Calculated MW	Observed MW
42.35 kDa	60-80 kDa

< 0.01 EU/μg of the protein by LAL method

Lyophilized from a 0.22  $\mu\text{m}$  filtered solution of PBS, pH 7.4.

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

This protein is a member of the neurotrophic tyrosine kinase receptor (NTRK) family. This kinase is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself and members of the MAPK pathway. The presence of this kinase leads to cell differentiation and may play a role in specifying sensory neuron subtypes. Mutations in this gene have been associated with congenital insensitivity to pain, anhidrosis, self-mutilating behavior, mental retardation and cancer. Alternate transcriptional splice variants of this gene have been found, but only three have been characterized to date.

Recombinant human Trk-A/NTRK1 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ala33-Glu413) of human Trk-A/NTRK1 (Accession #NP\_001012331.1) fused with and a 6×His tag at the C-terminus.

Measured by its ability to inhibit NGF-induced proliferation of TF-1 human erythroleukemic cells. The ED50 for this effect is 0.086-0.34 µg/mL in the presence of 10 ng/mL of Recombinant Human beta-NGF.


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.  
After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.  
Avoid repeated freeze/thaw cycles.

 | 400-999-6126


 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

# Validation Data



Recombinant human Trk-A/NTRK1 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Recombinant human Trk-A/NTRK1 inhibit NGF-induced proliferation of TF-1 human erythroleukemic cells. The ED50 for this effect is 0.086-0.34  $\mu$ g/mL in the presence of 10 ng/mL of Recombinant Human beta-NGF.