

Biotinylated Recombinant SARS-COV-2 Spike RBD Protein

Catalog No.: RP02326 Recombinant 1 Publications

Sequence Information

Species Gene ID Swiss Prot SARS-COV-2 QHD43416.1

Tags

C-His&Avi

Synonyms

S protein RBD;Spike glycoprotein Receptor-binding domain;S glycoprotein RBD;Spike protein RBD

Product Information

Source

Purification

HEK293 cells

> 95% by Tris-Bis PAGE;> 95% by SEC-HPLC

Endotoxin

< 1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
<u>~</u>	www.abclonal.com.cn

Background

Basic Information

Description

Biotinylated Recombinant SARS-COV-2 SARS-COV-2 Spike RBD Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Arg319-Asn532) of SARS-COV-2 SARS-COV-2 Spike RBD fused with His tag and Avi at the Cterminal.

Bio-Activity

Immobilized Human ACE2,His Tag at 2 μ g/mL (100 μ L/Well) on the plate. Dose response curve for Biotinylated SARS COV-2 Spike RBD,His Tag with the EC₅₀ of 0.097 μ g/mL determined by ELISA.


Storage

Store the lyophilized protein at -20°C to -80°C for long term.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data



Biotinylated SARS-COV-2 Spike RBD on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Biotinylated SARS-COV-2 Spike RBD is greater than 95% as determined by SEC-HPLC.



Immobilized Human ACE2, His Tag at $2\mu g/ml$ (100 μ l/Well) on the plate. Dose response curve for Biotinylated SARS COV-2 Spike RBD, His Tag with the EC $_{50}$ of $0.097\mu g/ml$ determined by ELISA.