

Recombinant SARS-COV-2 Spike S1+S2 ECD (S-ECD) Protein

Catalog No.: RP02334 Recombinant

Sequence Information

Species Gene ID Swiss Prot SARS-COV-2 43740568 OHD43416.1

Tags

C-hFc

Synonyms

S protein;Spike glycoprotein;S glycoprotein;COVID-19

Product Information

Source Purification

HEK293 cells \geq 95 % as

determined by Tris-Bis PAGE;≥ 95 % as determined by

HPLC.

Calculated MW Observed MW

158.3 kDa 230-250 kDa

Endotoxin

< 1 EU/ μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

8		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	Τ	www.abclonal.com.cn

Background

Basic Information

Description

Recombinant SARS-COV-2 Spike S1+S2 ECD (S-ECD) Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Val16-Glu1188) of SARS-COV-2 Spike S fused with hFc tag at the C-terminal.

Bio-Activity

Immobilized Human ACE2,His Tag at 5 μ g/mL (100 μ L/Well). Dose response curve for SARS-COV-2 Spike S,hFc Tag with the EC $_{50}$ of 0.9 μ g/mL determined by ELISA.


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data



Immobilized Human ACE2,His Tag at 5μ g/ml (100 μ l/Well). Dose response curve for SARS-COV-2 Spike S,hFc Tag with the EC $_{50}$ of 0.9 μ g/ml determined by ELISA.



Recombinant SARS-COV-2 Spike S1+S2 ECD (S-ECD) Protein was determined by Tris-Bis PAGE under reducing conditions.



The purity of SARS-COV-2 Spike S is greater than 95% as determined by SEC-HPLC.