

Biotinylated Recombinant Human Oncostatin-M/OSM Protein

Catalog No.: RP02464B Recombinant

Sequence Information

SpeciesGene IDSwiss ProtHuman5008P13725

Tags C-His&Avi

Synonyms OSM; oncostatin-M

Product Information

Source Purification

HEK293 cells ≥ 95 % as determined by Tris-Bis PAGE;≥ 95 % as determined by

HPLC.

Calculated MW Observed MW

25.1 kDa 30-40 kDa

Endotoxin

< 1 EU/ μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μ m filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

8		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	Τ	www.abclonal.com.cn

Background

OSM is a pleiotropic cytokine that initiates its biological activities by binding to specific cell surface receptors. Inhibits the proliferation of a number of tumor cell lines. Stimulates proliferation of AIDS-KS cells. It regulates cytokine production, including IL-6, G-CSF and GM-CSF from endothelial cells. Uses both type I OSM receptor (heterodimers composed of LIFR and IL6ST) and type II OSM receptor (heterodimers composed of OSMR and IL6ST). Involved in the maturation of fetal hepatocytes, thereby promoting liver development and regeneration.

Basic Information

Description

Biotinylated Recombinant Human Oncostatin-M/OSM Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Ala26-Arg221) of Human Oncostatin-M fused with a His tag and Avi tag at the C-terminal.

Bio-Activity

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80°C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.


Avoid repeated freeze/thaw cycles.

Operational Notes


For your safety and health, please wear a lab coat and disposable gloves for handling.

^{*} DODOODOODOODOODOO


Validation Data



Biotinylated Recombinant Human Oncostatin-M/OSM Protein was determined by Tris-Bis PAGE under reducing conditions.



The purity of Biotinylated Oncostatin M is greater than 95% as determined by SEC-HPLC.



Human LIF R, His Tag immobilized on CM5 Chip can bind Biotinylated Human Oncostatin M, His Tag with an affinity constant of 0.94 μM as determined in SPR assay (Biacore T200).