

Recombinant Cynomolgus B7-H1/PD-L1/CD274 Protein

Catalog No.: RP02470 Recombinant

Sequence Information

Species Gene ID Swiss Prot Cynomolgus 102145573 G7PSE7

Tags

C-His

Synonyms

CD274 antigenMGC142294; CD274 molecule; CD274; PDL1; PD-L1; PD-L1B7 homolog 1; B7-H; B7H1; B7-H1; B7H1PDCD1L1; PDCD1LG1; PDCD1LG1MGC142296; PDL1PDCD1 ligand 1; programmed cell death 1 ligand 1; Programmed death ligand 1; PDL1

Product Information

Source

HEK293 cells

≥ 95 % as determined by Tris-Bis PAGE;≥ 95 % as determined by

Purification

HPLC.

Calculated MW Observed MW

26 kDa 35-45 kDa

Endotoxin

< 1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

8

400-999-6126

Background

Basic Information

Description

Recombinant Cynomolgus PD-L1 Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Phe19-Arg238) of Cynomolgus PD-L1 fused with a His tag at the C-terminal.

Bio-Activity

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Validation Data




Recombinant Cynomolgus B7-H1/PD-L1/CD274 Protein was determined by Tris-Bis PAGE under reducing conditions.



The purity of Cynomolgus PD-L1 is greater than 95% as determined by SEC-HPLC.



Immobilized Cynomolgus PD-L1, His Tag at $0.5\mu g/ml$ (100 $\mu l/Well$) on the plate. Dose response curve for Anti-PD-L1 Antibody, hFc Tag with the EC₅₀ of 4.1ng/ml determined by ELISA.



Cynomolgus PD-1, hFc Tag captured on CM5 Chip via Protein A can bind Cynomolgus PD-L1, His Tag with an affinity constant of 1.28 μ M as determined in SPR assay (Biacore T200).