

Recombinant Cynomolgus Fc gamma RIIA/CD32a Protein

Catalog No.: RP02641 Recombinant

Sequence Information

Species Gene ID Swiss Prot Cynomolgus 102143568 Q8SPW4-1

Tags

C-His

Synonyms

CD32MGC23887; CDw32fc-gamma-Rlla; Fc gamma RllA; FCG2; Fc-gamma-Rlla; FcGR; FCGR2; FCGR2A; FCGR2A1; FcgRllA; FCRllA; fcRll-a; IGFR2MGC30032; IgG Fc receptor Il-a; FCG2: CD32A

Product Information

Source

Purification

HEK293 cells

≥ 95 % as determined by Tris-Bis PAGE;≥ 95 % as determined by HPLC.

Calculated MW Observed MW

21.3 kDa 30-40 kDa

Endotoxin

< 1 EU/ μg of the protein by LAL method.

Formulation

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
\odot	www.abclonal.com.cn

Background

The Fc gamma Rs have been divided into three classes based on close relationships in their extracellular domains; these groups are designated Fc gamma RI (also known as CD64), Fc gamma RII (CD32), and Fc gamma RIII (CD16). Each group may be encoded by multiple genes and exist in different isoforms depending on species and cell type. The CD64 proteins are high affinity receptors (~10e-8-10e-9 M) capable of binding monomeric IgG, whereas the CD16 and CD32 proteins bind IgG with lower affinities (~10e-6-10e-7 M) only recognizing IgG aggregates surrounding multivalent antigens.

Basic Information

Description

Recombinant Cynomolgus Fc gamma RIIA/CD32a Protein is expressed from Expi293 with His tag at the C-terminal. [] It contains Gln28-Pro208.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20 $^{\circ}$ C for 3 months, at 2-8 $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data



Recombinant Cynomolgus Fc gamma RIIA/CD32a Protein was determined by Tris-Bis PAGE under reducing conditions.



The purity of Cynomolgus Fc gamma RIIA is greater than 95% as determined by SEC-HPLC.



Rituximab captured on CM5 Chip via Protein A can bind Cynomolgus Fc gamma RIIA, His Tag with an affinity constant of 3.65 μM as determined in SPR assay (Biacore T200).