

Catalog No.: RP03107 **Recombinant**

Species	Gene ID	Swiss Prot
Human	4684	P13591-2

C-His

N-CAM-1; NCAM-1; NCAM1; CD56; NCAM;
MSK39

Source	Purification
HEK293 cells	≥ 95 % as determined by SDS-PAGE

78.23 kDa 100-140 kDa

< 0.01 EU/μg of the protein by LAL method

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn


Neural Cell Adhesion Molecule 1 (NCAM-1), a multifunctional member of the immunoglobulin superfamily, is expressed on the surface of neurons, glia, skeletal muscle, and natural killer cells. NCAM-1 has been implicated as having a role in cell-cell adhesion, involved in development of the nervous system, and for cells involved in the expansion of T cells and dendritic cells which play an important role in immune surveillance.

Recombinant Human NCAM-1/CD56 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Leu20-Gly718) of Human NCAM-1/CD56 (Accession #NP_851996.2) fused with a 6×His tag at the C-terminus.


Measured by the ability of the immobilized protein to support the adhesion of Neuro-2A Human neuroblastoma cells. The ED50 for this effect is 1.8-7.2 µg/mL.

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.
After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.
Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human NCAM-1/CD56 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



immobilized protein to support the adhesion of Neuro-2A Human neuroblastoma cells. The ED50 for this effect is 1.8-7.2 μ g/mL.