

Catalog No.: RP03391LQ **Recombinant**

Species	Gene ID	Swiss Prot
Human	1969	P29317

Tags
N-GST

EPHA2; ECK; Ephrin type-A receptor 2;
Epithelial cell kinase; ARCC2; CTPA;
CTPP1; CTRCT6

Source	Purification
Baculovirus-Insect Cells	≥ 90 % as determined by SDS-PAGE; ≥ 90 % as determined by HPLC.

Calculated MW	Observed MW
73.8 kDa	60-75 kDa

< 1 EU/μg of the protein by LAL method.

Supplied as a 0.22 μm filtered solution in 50 mM Tris-HCl, 150 mM NaCl, 5% glycerol, 5 mM DTT, 0.1 M Trehalose. (pH 7.5). Contact us for customized product form or formulation.

Please use running water to thaw it quickly.

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

EPH receptor A2 (ephrin type-A receptor 2) is a protein that in humans is encoded by the *EPHA2* gene. This gene belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family. EPH and EPH-related receptors have been implicated in mediating developmental events, particularly in the nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin receptors are divided into two groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. This gene encodes a protein that binds ephrin-A ligands.

Recombinant Human EphA2/ECK Kinase is produced by Baculovirus-Insect Cells expression system. The target protein is expressed with sequence (Arg561-Ile976) of Human EPHA2 (Accession #P29317) fused with a N-GST tag.

The activity of EPHA2 is based on the MSA technology, and the content and ratio of the substrate and the product are directly separated and detected in real time and dynamically by the different migration rates of the substrate and the product after the enzymatic reaction.


The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Store at -70°C. This product is stable at $\leq -70^{\circ}\text{C}$ for up to 1 year from the date of receipt. For optimal storage, aliquot into smaller quantities after centrifugation and store at recommended temperature.


Aliquots below 10 μ L are not advisable. Product must not be stored in diluted solutions. Avoid repeated freeze-thaw cycles.

For your safety and health, please wear a lab coat and disposable gloves for handling.

Validation Data



Recombinant Human EphA2/ECK Kinase was resolved with SDS-PAGE under reducing (Lane 1) and non-reducing (Lane 2) conditions.



The activity of EPHA2 is based on the MSA technology, and the content and ratio of the substrate and the product are directly separated and detected in real time and dynamically by the different migration rates of the substrate and the product after the enzymatic reaction.