

Catalog No.: RP03435LQ **Recombinant**

Species	Gene ID	Swiss Prot
Human	57787	O96L34

Synonyms
MARK4; KIAA1860; MARKL1;
MAP/microtubule affinity-regulating
kinase 4

Source E. coli	Purification ≥ 90 % as determined by SDS- PAGE; ≥ 90 % as determined by HPLC.
--------------------------	---

Calculated MW	Observed MW
39.5 kDa	30-40 kDa


< 1 EU/μg of the protein by LAL method.

Supplied as a 0.22 µm filtered solution in 50 mM Tris-HCl, 200 mM NaCl, 5% glycerol, 1 mM DTT. (pH 7.5). Contact us for customized product form or formulation.

Please use running water to thaw it quickly.

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

MAP/microtubule affinity-regulating kinase 4 is an enzyme that in humans is encoded by the MARK4 gene. MARK4 belongs to the family of serine/threonine kinases that phosphorylate microtubule-associated proteins (MAP) causing their detachment from microtubules. Detachment thereby increases microtubule dynamics and facilitates a number of cell activities including cell division, cell cycle control, cell polarity determination, and cell shape alterations. There are four members of the MARK protein family, MARK1-4, which are highly conserved. MARK4 kinase has been shown to be involved in microtubule organization in neuronal cells. Levels of MARK4 are elevated in Alzheimer's disease and may contribute to the pathological phosphorylation of tau protein in this disease.

Recombinant Human MARK4 Kinase is produced by E. coli expression system. The target protein is expressed with sequence (Asn44-Lys370) of Human MARK4 (Accession #Q96L34) fused with a N-His tag.


The activity of MARK4 is based on the ADP-GLO kinase activity assay quantifies kinase activity by measuring the conversion of ATP to ADP catalyzed by the kinase. Specific reagents are used to convert the ADP in the reaction back to ATP, resulting in the production of a luminescent signal.

Store at -70°C. This product is stable at $\leq -70^{\circ}\text{C}$ for up to 1 year from the date of receipt. For optimal storage, aliquot into smaller quantities after centrifugation and store at recommended temperature.


Aliquots below 10 μ L are not advisable. Product must not be stored in diluted solutions. Avoid repeated freeze-thaw cycles.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human MARK4 Kinase was resolved with SDS-PAGE under reducing (Lane 1) conditions.



The activity of MARK4 is based on the ADP-GLO kinase activity assay quantifies kinase activity by measuring the conversion of ATP to ADP catalyzed by the kinase. Specific reagents are used to convert the ADP in the reaction back to ATP, resulting in the production of a luminescent signal.