

Recombinant Human Syndecan-4 Protein

Catalog No.: RP03558 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	6385	P31431

Tags

C-His

Synonyms

Syndecan-4; SYND4; Amphiglycan;
Ryudocan core protein;SDC4

Product Information

Source	Purification
HEK293 Cells	> 95 % as determined by SDS-PAGE.

Calculated MW	Observed MW
15.3 kDa	25-35 kDa

Endotoxin

< 1.0 EU per µg protein as determined by the LAL method.

Formulation

Lyophilized from sterile PBS, pH 7.4.
Please contact us for any concerns or special requirements. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.

Reconstitution

Centrifuge the vial before opening.
Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Background

Syndecan-4 is a transmembrane heparan sulfate proteoglycan that co-operates with integrins during cell-matrix interactions for the assembly of focal adhesions and actin stress fibers and in the phosphorylation of focal adhesion kinase (FAK) on Tyr397. Syndecan-4 plays roles in the formation of focal adhesions and stress fibers. The cytoplasmic domain of syndecan-4 interacts with several signalling and structural proteins, and both extracellular and cytoplasmic domains are necessary for regulated activation of associated transmembrane receptors. Syndecan-4/SDC4 is a heparan sulfate proteoglycan and works as a coreceptor for various growth factors. SDC4 deficiency limits neointimal formation after vascular injury by regulating vascular smooth muscle cells (VSMCs) proliferation and vascular progenitor cells (VPCs) mobilization. Therefore, SDC4 may be a novel therapeutic target for preventing arterial restenosis after angioplasty.

Basic Information

Description

Recombinant Human Syndecan-4 Protein is produced by HEK293 cells system. The target protein is expressed with sequence (Met1-Glu145) of Human Syndecan-4 (Accession #NP_002990.2) fused with His at the C-Terminus.

Bio-Activity

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Contact

 | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Validation Data



Recombinant Human Syndecan-4 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.